

SUPPLEMENTARY MATERIAL TO
Modelling and optimisation of activated sludge process using artificial neural networks and genetic algorithms

SAURABH SAHADEV^{1*}, GOPAL MADHU² and ROY M. THOMAS²

¹Department of Chemical Engineering, Government Engineering College, Thrissur, Kerala, India and ²School of Engineering, Cochin University of Science and Technology, Kochi, Kerala, India

J. Serb. Chem. Soc. 91 (0) (2026) 000–000




Fig. S-1. Relationship between pH_{Inf} and pH_{Eff}




Fig. S-2. Relationship between SS_{Inf} and pH_{Eff}

* Corresponding author. E-mail: saurabhsdev@gmail.com




Fig. S-3. Relationship between BOD_{inf} and pH_{eff}




Fig. S-4. Relationship between $O\&G_{inf}$ and pH_{eff}




Fig. S-5. Relationship between pH_{inf} and SS_{eff}




Fig. S-6. Relationship between SS_{inf} and SS_{eff}




Fig. S-7. Relationship between BOD_{Inf} and SS_{Eff}




Fig. S-8. Relationship between $O\&G_{Inf}$ and SS_{Eff}




Fig. S-9. Relationship between pH_{Inf} and BOD_{Eff}




Fig. S-10. Relationship between SS_{Inf} and BOD_{Eff}




Fig. S-11. Relationship between BOD_{Inf} and BOD_{Eff}




Fig. S-12. Relationship between $O\&G_{Inf}$ and BOD_{Eff}