



SUPPLEMENTARY MATERIAL TO
**Magnetic *Ambrosia artemisiifolia*-based biosorbent for Congo red
removal from water: A circular economy approach**

NATALIJA NEDIĆ¹, TAMARA TADIĆ¹, BOJANA MARKOVIĆ¹, ALEKSANDRA
NASTASOVIĆ¹, ALEKSANDAR POPOVIĆ², AND SANDRA BULATOVIĆ^{1*}

¹Institute of Chemistry Technology and Metallurgy, University of Belgrade, Njegoševa 12, 11000
Beograd, Serbia, and ²University of Belgrade, Faculty of Chemistry, Studentski trg 12–16,
11158 Belgrade, Serbia.

RESULTS AND DISCUSSION

Biosorption kinetics

Table SI. Kinetic models applied to analyze CR biosorption with Fe₃O₄/MB.¹

Model	Equation	Plot	Parameter
Pseudo-first-order (PFO)	$\log(Q_e - Q_t) = \log(Q_e) - \frac{k_1}{2.303}t$	$\log(Q_e - Q_t)$ vs. t	Q_e, k_1
Pseudo-second-order (PSO)	$\frac{t}{Q_t} = \frac{1}{k_2 Q_e^2} + \frac{t}{Q_e}$	$\frac{t}{Q_t}$ vs. t	Q_e, k_2
Elovich	$Q_t = \frac{1}{\beta} \ln(\alpha\beta) + \frac{1}{\beta} \ln(t)$	Q_t vs. $\ln(t)$	α, β
Intra-particle diffusion (IPD)	$Q_t = k_3 t^{0.5} + C_{id}$	Q_t vs. $t^{0.5}$	k_3

where Q_e and Q_t (mg g⁻¹) are the adsorption capacity's at equilibrium and at time t , k_1 (min⁻¹) and k_2 (g mg⁻¹ min⁻¹) are the rate constant; α (mg g⁻¹) is the initial adsorption rate, β (g mg⁻¹) is related to the extent of surface coverage; k_3 (g mg⁻¹ min⁻¹) is intra-particle diffusion rate constant

* Corresponding author. E-mail: sandra.bulatovic@ihtm.bg.ac.rs

*Biosorption isotherms***Table SII.** Isotherm models applied to analyze CR biosorption with Fe₃O₄/MB.²

Model	Equation	Plot	Parameter
Langmuir	$\frac{C_e}{Q_e} = \frac{1}{k_L Q_{\max}} + \frac{C_e}{Q_{\max}}$	$\frac{C_e}{Q_e}$ vs. C_e	$Q_{\max}, k_L$
Freundlich	$\log Q_e = \log k_F + \frac{1}{n} \log C_e$	$\log Q_e$ vs. $\log C_e$	k_F, n
Temkin	$Q_e = \frac{RT}{b_T} \ln A_T + \frac{RT}{b_T} \ln C_e$	Q_e vs. $\ln C_e$	A_T, b_T
Elovich	$\ln \frac{Q_e}{C_e} = \ln K_E Q_m - \frac{Q_e}{Q_m}$	$\ln \frac{Q_e}{C_e}$ vs. Q_e	Q_m, K_E

where Q_e (mg g⁻¹) and C_e (mg L⁻¹) are the adsorption capacity at equilibrium and adsorbate equilibrium concentration respectively; k_L (L mg⁻¹) is Langmuir constant; Q_m (mg g⁻¹) is the maximum monolayer adsorption capacity; n and k_F (L g⁻¹) are Freundlich constants related to adsorption and intensity of adsorption driving force or surface heterogeneity, respectively; A_T is Temkin constants (L mg⁻¹), T is the temperature (K), R is the gas constant (8.314 J mol⁻¹ K⁻¹); K_E (L g⁻¹) is Elovich equilibrium constant.

REFERENCES

1. M. A. Fawzy, M. Gomaa, *J. Environ. Manage.* **262** (2020) 110380 (<https://doi.org/10.1016/j.jenvman.2020.110380>)
2. A. N. El-hoshoudy, M. Matallah, H. Gouzi, B. Saidat, Y. Khane, M. Chabani, I. Nouacer, *Int. J. Environ. Sci. Technol.* **21** (2024) 10187 (<https://doi.org/10.1007/s13762-024-05616-6>).