

SUPPLEMENTARY MATERIAL TO  
**A novel carbon paste electrode based on nitrogen-doped hydrothermal carbon for electrochemical determination of carbendazim**

ANA KALIJADIS<sup>1\*</sup>, JELENA ĐORĐEVIĆ<sup>1</sup>, ZSIGMOND PAPP<sup>2</sup>, BOJAN JOKIĆ<sup>3</sup>,  
VUK SPASOJEVIĆ<sup>1</sup>, BILJANA BABIĆ<sup>1</sup> and TATJANA TRTIĆ-PETROVIĆ<sup>1</sup>

<sup>1</sup>Vinča Institute of Nuclear Sciences, University of Belgrade, P. O. Box 522, 11001 Belgrade,

<sup>2</sup>Faculty of Biofarming, John Naisbitt University, Maršala Tita 39, 24300 Bačka

Topola, Serbia and <sup>3</sup>Faculty of Technology and Metallurgy, University of Belgrade,  
Karnegijeva 4, 11000 Belgrade, Serbia

J. Serb. Chem. Soc. 82 (11) (2017) 1259–1272




Figure S-1. Effect of TCP mass fraction in CPE/CHTCN<sub>0.1</sub> on cyclic voltammograms of Fe(CN)<sub>6</sub><sup>3-/4-</sup> redox couple in 0.1 M K<sub>2</sub>SO<sub>4</sub>.

\*Corresponding author. E-mail: anaudovicic@vin.bg.ac.rs