

SUPPLEMENTARY MATERIAL TO

Expression, purification and characterization of cellobiose dehydrogenase mutants from *Phanerochaete chrysosporium* in *Pichia pastoris* KM71H strain

ANA MARIJA J. BALAŽ¹, MARIJA B. BLAŽIĆ¹, NIKOLINA POPOVIĆ², OLIVERA L.
PRODANOVIĆ³, RALUCA V. OSTAFE⁴, RAINER FISCHER⁵ and RADIVOJE M.
PRODANOVIĆ^{2*}

¹*Institute of Chemistry, Technology and Metallurgy, University of Belgrade, Njegoseva 12, 11000 Belgrade, Serbia*

²Faculty of Chemistry, University of Belgrade, Studentski trg 12 – 16, 11000 Belgrade, Serbia

³ Institute for Multidisciplinary Studies, University of Belgrade, Kneza Višeslava 1, 11030 Belgrade, Serbia

⁴ Molecular Evolution Protein Engineering and Production facility (MEPEP), Purdue University,
207 S. Martin Jischke Dr., West Lafayette, IN 47907, USA

⁵*Indiana Bioscience Research Institute, Single Cell Analytics Center, 1345 W. 16th St. Suite 300,
Indianapolis, IN 46202, USA*

*Corresponding author. E-mail: rprodano@chem.bg.ac.rs

28


29

30

31 TABLE S1. Primers used for creation of triple mutant and error prone library mutants using
 32 wtCDH – pPICZ α A vector as template


Primer name	Primer sequence
Forward primer D20N	GGTATCACCAACCCTGTTCATG
Forward primer A64T	CTCGGTGGCACCATGAACAAC
Forward primer V592M	CGCAGCCTCCATGAACCTCC
Forward primer D20N – V22A	CACCAACCCTGCTCATGACG
Forward primer T84A	TTTCCTCCGCTCGCTGG
Forward primer A261P	ACGTATGTCCCTCCATG
Forward primer E674G	AGACACTCGGGAGTACG
Forward primer N715S	TTGGCACGAGCAACCTGTT

33



35 Fig. S1. Ion-exchange chromatography of tm CDH


36



38 Fig. S2. Ion-exchange chromatography of H5 CDH

39

40




41

42

Fig. S3. Ion-exchange chromatography of H9 CDH

43



44

45

Fig. S4. Gel filtration of tm CDH

46




Fig. S5. Gel filtration of H5 CDH

50

51

52




Fig. S6. Gel filtration of H9 CDH

53 TABLE S2 Purification table for CDH. FM-fermentation media, UF-ultrafiltrate, DEAE-sample
 54 after ion-exchange chromatography, GF-sample after gel filtration. *Cp*-protein concentration, *Ac*-
 55 enzyme activity, *V*-volume, *Y*-yield of purification, *Pf*-purification factor, *Sp*-specific enzyme
 56 activity.

	tm CDH				H5 CDH				H9 CDH			
	FM	UF	DEAE	GF	FM	UF	DEAE	GF	FM	UF	DEAE	GF
<i>Cp</i> / mg mL ⁻¹	0.59	0.42	0.48	0.08	0.52	0.45	0.74	0.17	0.46	0.20	0.15	0.024
<i>Ac</i> / IU mL ⁻¹	0.91	2.34	5.41	1.17	1.99	7.30	16.5	4.89	0.42	1.10	1.87	0.34
<i>V</i> / mL	50	12	3	7	50	12	3	6	50	12	3	5
<i>Y</i> / %	100	62	36	18	100	88	50	29	100	63	27	8.2
<i>Pf</i>	1	3.6	7.2	9.1	1	4.2	5.8	7.3	1	5.9	13	16
<i>Sp</i> / IU mg ⁻¹	1.55	5.55	11.2	14.1	3.83	16.2	22.4	28.1	0.92	5.44	12.2	14.5

57

58

59